

HAIR ELEMENTS



LAB#:
 PATIENT:
 SEX: Female
 AGE: 35
 CLIENT#:

POTENTIALLY TOXIC ELEMENTS				
TOXIC ELEMENTS	RESULT µg/g	REFERENCE RANGE	PERCENTILE	
			68 th	95 th
Aluminum	7.8	< 7.0	████████████████████	████████████████████
Antimony	0.010	< 0.050	●	████████████████████
Arsenic	0.074	< 0.060	████████████████████	████████████████████
Beryllium	< 0.01	< 0.020	████████████████████	████████████████████
Bismuth	0.081	< 0.10	████████████████████	████████████████████
Cadmium	0.013	< 0.10	●	████████████████████
Lead	0.33	< 1.0	████████████████████	████████████████████
Mercury	1.3	< 1.1	████████████████████	████████████████████
Platinum	< 0.003	< 0.005	████████████████████	████████████████████
Thallium	< 0.001	< 0.010	████████████████████	████████████████████
Thorium	< 0.001	< 0.005	████████████████████	████████████████████
Uranium	< 0.001	< 0.060	████████████████████	████████████████████
Nickel	0.29	< 0.40	████████████████████	████████████████████
Silver	0.02	< 0.15	████████████████████	████████████████████
Tin	0.09	< 0.30	████████████████████	████████████████████
Titanium	4.5	< 1.0	████████████████████	████████████████████
Total Toxic Representation			████████████████████	████████████████████

ESSENTIAL AND OTHER ELEMENTS							
ELEMENTS	RESULT µg/g	REFERENCE RANGE	PERCENTILE				
			2.5 th	16 th	50 th	84 th	97.5 th
Calcium	1300	300- 1200	████████████████████	████████████████████	████████████████████	████████████████████	████████████████████
Magnesium	100	35- 120	████████████████████	████████████████████	████████████████████	████████████████████	████████████████████
Sodium	55	12- 90	████████████████████	████████████████████	████████████████████	████████████████████	████████████████████
Potassium	130	8- 38	████████████████████	████████████████████	████████████████████	████████████████████	████████████████████
Copper	14	12- 35	████████████████████	████████████████████	████████████████████	████████████████████	████████████████████
Zinc	220	140- 220	████████████████████	████████████████████	████████████████████	████████████████████	████████████████████
Manganese	0.18	0.15- 0.65	████████████████████	████████████████████	████████████████████	████████████████████	████████████████████
Chromium	0.36	0.20- 0.40	████████████████████	████████████████████	████████████████████	████████████████████	████████████████████
Vanadium	0.19	0.018- 0.065	████████████████████	████████████████████	████████████████████	████████████████████	████████████████████
Molybdenum	0.041	0.028- 0.056	████████████████████	████████████████████	████████████████████	████████████████████	████████████████████
Boron	0.67	0.30- 2.0	████████████████████	████████████████████	████████████████████	████████████████████	████████████████████
Iodine	0.17	0.25- 1.3	████████████████████	████████████████████	████████████████████	████████████████████	████████████████████
Lithium	0.012	0.007- 0.023	████████████████████	████████████████████	████████████████████	████████████████████	████████████████████
Phosphorus	225	160- 250	████████████████████	████████████████████	████████████████████	████████████████████	████████████████████
Selenium	0.58	0.95- 1.7	████████████████████	████████████████████	████████████████████	████████████████████	████████████████████
Strontium	2.3	0.50- 7.6	████████████████████	████████████████████	████████████████████	████████████████████	████████████████████
Sulfur	49000	44500- 52000	████████████████████	████████████████████	████████████████████	████████████████████	████████████████████
Barium	0.61	0.26- 3.0	████████████████████	████████████████████	████████████████████	████████████████████	████████████████████
Cobalt	0.006	0.013- 0.050	████████████████████	████████████████████	████████████████████	████████████████████	████████████████████
Iron	6.7	5.4- 14	████████████████████	████████████████████	████████████████████	████████████████████	████████████████████
Germanium	0.034	0.045- 0.065	████████████████████	████████████████████	████████████████████	████████████████████	████████████████████
Rubidium	0.14	0.007- 0.096	████████████████████	████████████████████	████████████████████	████████████████████	████████████████████
Zirconium	0.060	0.020- 0.42	████████████████████	████████████████████	████████████████████	████████████████████	████████████████████

SPECIMEN DATA			RATIOS		
COMMENTS: 98443			ELEMENTS	RATIOS	EXPECTED RANGE
Date Collected: 7/20/2006	Sample Size: 0.201 g		Ca/Mg	13	4- 30
Date Received: 7/31/2006	Sample Type: Head		Ca/P	5.78	1- 12
Date Completed: 8/2/2006	Hair Color:		Na/K	0.423	0.5- 10
Methodology: ICP-MS	Treatment:		Zn/Cu	15.7	4- 20
	Shampoo:		Zn/Cd	> 999	> 800

V06.99

The Great Plains Laboratory, Inc. • 11813 W. 77 Street, Lenexa KS, 66214 • Tel: 913.341.8949 • Fax: 913.341.6207

Analyzed by ©DOCTOR'S DATA, INC. • ADDRESS: 3755 Illinois Avenue, St. Charles, IL 60174-2420 • CLIA ID NO: 14D0646470 • MEDICARE PROVIDER NO: 148453

Lab number:
Patient:

Hair

Page: 1
Client:

HAIR ELEMENTS REPORT INTRODUCTION

Hair is an excretory tissue for essential, nonessential and potentially toxic elements. In general, the amount of an element that is irreversibly incorporated into growing hair is proportional to the level of the element in other body tissues. Therefore, hair elements analysis provides an indirect screening test for physiological excess, deficiency or maldistribution of elements in the body. Clinical research indicates that hair levels of specific elements, particularly potentially toxic elements such as cadmium, mercury, lead and arsenic, are highly correlated with pathological disorders. For such elements, levels in hair may be more indicative of body stores than the levels in blood and urine.

All screening tests have limitations that must be taken into consideration. The correlation between hair element levels and physiological disorders is determined by numerous factors. Individual variability and compensatory mechanisms are major factors that affect the relationship between the distribution of elements in hair and symptoms and pathological conditions. It is also very important to keep in mind that scalp hair is vulnerable to external contamination of elements by exposure to hair treatments and products. Likewise, some hair treatments (e.g. permanent solutions, dyes, and bleach) can strip hair of endogenously acquired elements and result in false low values. Careful consideration of the limitations must be made in the interpretation of results of hair analysis. The data provided should be considered in conjunction with symptomology, diet analysis, occupation and lifestyle, physical examination and the results of other analytical laboratory tests.

Caution: The contents of this report are not intended to be diagnostic and the physician using this information is cautioned against treatment based solely on the results of this screening test. For example, copper supplementation based upon a result of low hair copper is contraindicated in patients afflicted with Wilson's Disease.

Aluminum High

The Aluminum (Al) level in hair is a reliable indicator of assimilation of this element, provided that hair preparations have not added exogenous Al. Al is a nonessential element that can be toxic if excessively assimilated into cells.

Excess Al can inhibit the formation of alpha-keto glutarate and result in toxic levels of ammonia in tissues. Al can bond to phosphorylated bases on DNA and disrupt protein synthesis and catabolism. Al excess should be considered when symptoms of presenile dementia or Alzheimer's disease are observed. Hair Al is commonly elevated in children and adults with low zinc and behavioral/learning disorders such as ADD, ADHD and autism. Individuals with renal problems or on renal dialysis may have elevated Al.

Possible sources of Al include some antacid medications, Al cookware, baking powder, processed cheese, drinking water, and antiperspirant components that may be absorbed. Analyses performed at DDI indicate extremely high levels of Al are in many colloidal mineral products.

Al has neurotoxic effects at high levels, but low levels of accumulation may not elicit immediate symptoms. Early symptoms of Al burden may include: fatigue, headache, and symptoms of phosphate depletion.

Lab number:
Patient:

Hair

Page: 2
Client:

A urine elements test can be used to corroborate Al exposure. Al can be effectively complexed and excreted with silicon (J. Environ. Pathol. Toxicol. Oncol., 13(3): 205-7, 1994). A complex of malic acid and Mg has been reported to be quite effective in lowering Al levels (DDI clients).

Arsenic High

In general, hair provides a rough estimate of exposure to Arsenic (As) absorbed from food and water. However, hair can be contaminated externally with As from air, water, dust, shampoos and soap. Inorganic As, and some organic As compounds, can cause toxicity. Some research suggests that As may be essential at extremely low levels but its function is not understood. Inorganic As accumulates in hair, nails, skin, thyroid gland, bone and the gastrointestinal tract. Organic As is rapidly excreted in the urine.

As can cause malaise, muscle weakness, vomiting, diarrhea, dermatitis, and skin cancer. Long-term exposure may affect the peripheral nervous, cardiovascular and hematopoietic systems. As is a major biological antagonist to selenium.

Common sources of As are insecticides (calcium and lead arsenate), well water, smog, shellfish (arsenobetaine), and industrial exposure, particularly in the manufacture of electronic components (gallium arsenide).

As burden can be confirmed by urine elements analysis. Comparison of urine As levels pre and post provocation (DMPS, DMSA, D-penicillamine) permit differentiation between recent uptake and body stores.

Mercury High

Mercury (Hg) is toxic to humans and animals. The accumulation of Hg in the body is generally reflected by the hair Hg levels, but hair Hg levels can be artifactually high in association with the use of certain hair dyes. Individuals vary greatly in sensitivity and tolerance to Hg burden.

At hair levels below 3 µg/g, Hg can suppress biological selenium function and may cause or contribute to immune dysregulation in sensitive individuals. Hallmark symptoms of excess Hg include: loss of appetite, decreased senses of touch, hearing, and vision, fatigue, depression, emotional instability, peripheral numbness and tremors, poor memory and cognitive dysfunction, and neuromuscular disorders. Hair Hg has been reported to correlate with acute myocardial infarction and on average each 1 µg/g of hair Hg was found to correlate with a 9% increase in AMI risk (Circulation 1995; 91:645-655).

Sources of Hg include dental amalgams, contaminated seafood, water supplies, some hemorrhoidal preparations, skin lightening agents, instruments (thermometers, electrodes, batteries), and combustion of fossil fuels, some fertilizers, and the paper/pulp and gold industries. After dental amalgams are installed or removed a transient (several months) increase in hair Hg is observed. Also, "baseline" hair Hg levels for individuals with dental amalgams are higher (about 1 to 2 µg/g) than are baseline levels for those without (below 1 µg/g).

Confirmatory tests for elevated Hg are measurement of whole blood as an indication of recent/ongoing exposure (does not correlate with whole body accumulation) and measurement of

Lab number:
Patient:

Hair

Page: 3
Client:

urine Hg following use of a dithiol chelating or mobilizing agent such as DMSA or DMPS (an indication of total body burden).

Potassium High

High hair Potassium (K) is not necessarily reflective of dietary intake or nutrient status. However, elevated K may be reflective of metabolic disorders associated with exposure to potentially toxic elements.

K is an electrolyte and a potentiator of enzyme functions, but neither of these functions take place in hair. Elevated K in hair may reflect overall retention of K by the body or maldistribution of this element. In adrenocortical insufficiency, K is increased in blood, while it is decreased in urine; cellular K may or may not be increased. Also, hair is occasionally contaminated with K from some shampoos. Observations at DDI indicate that K and sodium levels in hair are commonly high in association with toxic element burden. The elevated K and sodium levels are often concomitant with low levels of calcium and magnesium in hair. This apparent phenomena requires further investigation.

Elevated hair potassium should be viewed as a screening test. Appropriate tests for excess body K include measurements of packed red blood cell K; serum or whole blood K and sodium/K ratio, measurement of urine K and sodium/K ratio; and an assessment of adrenocortical function.

Copper Normal

Hair Copper (Cu) levels are usually indicative of body status, except that exogenous contamination may occur giving a false normal (or false high). Common sources of contamination include: permanent solutions, dyes, bleaches, and swimming pools/hot tubs in which Cu compounds have been used as algacides.

Cu is an essential element that activates specific enzymes. Erythrocyte superoxide dismutase (SOD) is a Cu (and zinc) dependent enzyme; lysyl oxidase which catalyzes crosslinking of collagen is another Cu dependent enzyme. Adrenal catecholamine synthesis is Cu dependent, because the enzyme dopamine beta-hydroxylase, which catalyzes formation of norepinephrine from dopamine, requires Cu.

If hair Cu is in the normal range, this usually means tissue levels are in the normal range. However, under circumstances of contamination, a real Cu deficit could appear as a (false) normal. If symptoms of Cu deficiency are present, a whole blood or red blood cell elements analysis can be performed for confirmation of Cu status.

Cobalt Low

One can not determine vitamin B-12 status by use of hair analysis, and the clinical significance of low hair Cobalt (Co) levels is not known. Hair is analyzed for Co primarily for detection of excessive intake of the potentially toxic element.

There is little evidence that Co has an essential function in humans other than as an obligatory constituent of the vitamin B-12 molecule. Humans absorb Co as inorganic Co and as vitamin B-12; the body pools of each fluctuate independently. Humans cannot convert inorganic Co to vitamin B-12.

Lab number:
Patient:

Hair

Page: 4
Client:

The dietary content of Co is highly variable, depending upon types of foods eaten, geographical location and type of soil. Vegetarians often have lower Co levels than meat eaters.

Appropriate tests for determination of vitamin B-12 status are the measurement of urine levels of methylmalonic acid (elevated with vitamin B-12 coenzyme deficiency/dysfunction), a quantitative blood assay for vitamin B-12, a urine amino acids analysis (several metabolic steps require vitamin B-12), and diet analysis.

Vanadium High

High levels of Vanadium (V) in hair may be indicative of excess absorption of the element. It is well established that excess V can have toxic effects in humans. Although it appears that V may have essential functions, over zealous supplementation is not warranted.

Excess levels of V in the body can result from chronic consumption of fish, shrimp, crabs, and oysters derived from water near offshore oil rigs (Metals in Clinical and Analytical Chemistry, 1994). Industrial/environmental sources of V include: processing of mineral ores, phosphate fertilizers, combustion of oil and coal, production of steel, and chemicals used in the fixation of dyes and print.

Symptoms of V toxicity vary with chemical form and route of absorption. Inhalation of excess V may produce respiratory irritation and bronchitis. Excess ingestion of V can result in decreased appetite, depressed growth, diarrhea/gastrointestinal disturbances, nephrotoxic and hematotoxic effects. Pallor, diarrhea, and green tongue are early signs of excess V and have been reported in human subjects consuming about 20 mg V/day (Modern Nutrition in Health and Disease, 8th edition, eds. Shils, M., Olson, J., and Mosha, S., 1994).

Confirmatory tests for excess V are red blood cell elements analysis, and urine V which reflects recent intake.

Selenium Low

Selenium (Se) is normally found in hair at very low levels, and several studies provide evidence that low hair Se is reflective of dietary intake and associated with cardiovascular disorders. Utilization of hair Se levels to assess nutritional status, however, is complicated by the fact that use of Se- or sulfur-containing shampoo markedly increases hair Se (externally) and can give a false high value.

Se is an extremely important essential element due to its antioxidative function as an obligatory component of the enzyme glutathione peroxidase. Se is also protective in its capacity to bind and "inactivate" mercury, and Se is an essential cofactor in the deiodination of T-4 to active T-3 (thyroid hormone). Some conditions of functional hypothyroidism therefore may be due to Se deficiency (Nature; 349:438-440, 1991); this is of particular concern with mercury exposure. Studies have also indicated significant inverse correlations between Se and heart disease, cancer, and asthma.

Selenium deficiency is common and can result from low dietary intake of Se or vitamin E, and exposure to toxic metals, pesticides/herbicides and chemical solvents.

Lab number:
Patient:

Hair

Page: 5
Client:

Symptoms of Se deficiency are similar to that of vitamin E deficiency and include muscle aches, increased inflammatory response, loss of body weight, alopecia, listlessness, skeletal and muscular degeneration, growth stunting, and depressed immune function.

Confirmatory tests for Se deficiency are Se content of packed red blood cells, and activity of glutathione peroxidase in red blood cells.

Titanium High

Titanium (Ti) is measured in hair to assist in the identification of external contamination of hair by treatments and products. Shampoos, dyes, and "highlighting" are the primary sources of Ti, which binds tenaciously to hair.

Ti dioxide is the most common form of Ti used for industrial purposes; e.g. coating of welding rods and as white pigment in paints, dyes, and paper fillers. Ti dioxide and other Ti containing compounds have extremely low toxicity. The elevated level of Ti in the hair sample is most likely without clinical significance.

Total Toxic Element Indication

The potentially toxic elements vary considerably with respect to their relative toxicities. The accumulation of more than one of the most toxic elements may have synergistic adverse effects, even if the level of each individual element is not strikingly high. Therefore, we present a total toxic element "score" which is estimated using a weighted average based upon relative toxicity. For example, the combined presence of lead and mercury will give a higher total score than that of the combination of silver and beryllium.

Lab number:
Patient:

Hair

Page: 6
Client:
